

## Mathematics C3

1. (a)
- |  |   |                 |                            |
|--|---|-----------------|----------------------------|
|  | 0   | 1               |                            |
|  | $\pi/12$  | $0.933012701$   |                            |
|  | $\pi/6$   | $0.75$          |                            |
|  | $\pi/4$   | $0.5$           | (5 values correct) B2      |
|  | $\pi/3$   | $0.25$          | (3 or 4 values correct) B1 |
|  | Correct formula with $h = \pi/12$   |                 | M1                         |
|  | $I \approx \frac{\pi/12}{3} \times \{1 + 0.25 + 4(0.933012701 + 0.5) + 2(0.75)\}$ |                 |                            |
|  | $I \approx 8.482050804 \times (\pi/12) \div 3$                                    |                 |                            |
|  | $I \approx 0.740198569$   |                 |                            |
|  | $I \approx 0.7402$  | (f.t. one slip) | A1                         |

**Note: Answer only with no working shown earns 0 marks**

- (b)
- |  |  |                                     |
|--|--|-------------------------------------|
|  | $\int_0^{\pi/3} \sin^2 x \, dx = \int_0^{\pi/3} 1 \, dx - \int_0^{\pi/3} \cos^2 x \, dx$ | M1                                  |
|  | $\int_0^{\pi/3} \sin^2 x \, dx = 0.3070$   | (f.t. candidate's answer to (a)) A1 |

**Note: Answer only with no working shown earns 0 marks**

2. (a)
- |  |  |    |
|--|--|----|
|  | e.g. $\theta = \pi/2, \phi = \pi$  |    |
|  | $\sin(\theta + \phi) = -1$ (choice of $\theta, \phi$ and one correct evaluation) | B1 |
|  | $\sin \theta + \sin \phi = 1$ (both evaluations correct but different)           | B1 |
- (b)
- |  |  |             |
|--|--|-------------|
|  | $\sec^2 \theta + 8 = 4(\sec^2 \theta - 1) + 5 \sec \theta$   |             |
|  | (correct use of $\tan^2 \theta = \sec^2 \theta - 1$ )  | M1          |
|  | An attempt to collect terms, form and solve quadratic equation in $\sec \theta$ , either by using the quadratic formula or by getting the expression into the form $(a \sec \theta + b)(c \sec \theta + d)$ , with $a \times c =$ candidate's coefficient of $\sec^2 \theta$ and $b \times d =$ candidate's constant |             |
|  | $3 \sec^2 \theta + 5 \sec \theta - 12 = 0 \Rightarrow (3 \sec \theta - 4)(\sec \theta + 3) = 0$  | m1          |
|  | $\Rightarrow \sec \theta = \frac{4}{3}, \sec \theta = -3$  |             |
|  | $\Rightarrow \cos \theta = \frac{3}{4}, \cos \theta = -\frac{1}{3}$  | (c.a.o.) A1 |
|  | $\theta = 41.41^\circ, 318.59^\circ$   |             |
|  | B1   |             |
|  | $\theta = 109.47^\circ, 250.53^\circ$  | B1 B1       |

**Note:** Subtract 1 mark for each additional root in range for each branch, ignore roots outside range.

$\cos \theta = +, -$ , f.t. for 3 marks,  $\cos \theta = -, -$ , f.t. for 2 marks  
 $\cos \theta = +, +$ , f.t. for 1 mark

3. (a) (i) candidate's  $x$ -derivative =  $6t$ ,  
candidate's  $y$ -derivative =  $6t^5 - 12t^2$   
(at least two of the three terms correct) B1  
 $\frac{dy}{dx} = \frac{\text{candidate's } y\text{-derivative}}{\text{candidate's } x\text{-derivative}}$  M1  
 $\frac{dy}{dx} = \frac{6t^5 - 12t^2}{6t}$  (c.a.o.) A1
- (ii)  $\frac{6t^5 - 12t^2}{6t} = \frac{7}{2}$  (f.t. candidate's expression from (i)) M1  
 $2t^4 - 4t - 7 = 0$   
(convincing) A1
- (b)  $f(t) = 2t^4 - 4t - 7$   
An attempt to check values or signs of  $f(t)$  at  $t = 1, t = 2$  M1  
 $f(1) = -9 < 0, f(2) = 17 > 0$   
Change of sign  $\Rightarrow f(t) = 0$  has root in (1, 2) A1  
 $t_0 = 1.6$   
 $t_1 = 1.608861654$  ( $t_1$  correct, at least 5 places after the point) B1  
 $t_2 = 1.609924568$   
 $t_3 = 1.610051919$   
 $t_4 = 1.610067175 = 1.61007$  ( $t_4$  correct to 5 decimal places) B1  
An attempt to check values or signs of  $f(t)$  at  $t = 1.610065,$   
 $t = 1.610075$  M1  
 $f(1.610065) = -1.25 \times 10^{-4} < 0, f(1.610075) = 1.69 \times 10^{-4} > 0$  A1  
Change of sign  $\Rightarrow \alpha = 1.61007$  correct to five decimal places A1

**Note: 'Change of sign' must appear at least once.**

4.  $\frac{d(x^2y^2)}{dx} = x^2 \times 2y \frac{dy}{dx} + 2x \times y^2$  B1  
 $\frac{d(2y^3)}{dx} = 6y^2 \times \frac{dy}{dx}$  B1  
 $\frac{d(x^4 - 2x + 6)}{dx} = 4x^3 - 2$  B1  
 $x = 2, y = 3 \Rightarrow \frac{dy}{dx} = \frac{66}{5} = \underline{11}$  (o.e.) (c.a.o.) B1

5. (a)  $\frac{dy}{dx} = \frac{4}{1 + (4x)^2}$  or  $\frac{1}{1 + (4x)^2}$  or  $\frac{4}{1 + 4x^2}$  M1  
 $\frac{dy}{dx} = \frac{4}{1 + 16x^2}$  A1
- (b)  $\frac{dy}{dx} = e^{x^3} \times f(x)$  ( $f(x) \neq 1$ ) M1  
 $\frac{dy}{dx} = 3x^2 \times e^{x^3}$  A1
- (c)  $\frac{dy}{dx} = x^5 \times f(x) + \ln x \times g(x)$  ( $f(x), g(x) \neq 1$ ) M1  
 $\frac{dy}{dx} = x^5 \times f(x) + \ln x \times g(x)$  (either  $f(x) = 1/x$  or  $g(x) = 5x^4$ ) A1  
 $\frac{dy}{dx} = x^4 + 5x^4 \times \ln x$  (c.a.o.) A1
- (d)  $\frac{dy}{dx} = \frac{(5 - 4x^2) \times f(x) - (3 - 2x^2) \times g(x)}{(5 - 4x^2)^2}$  ( $f(x), g(x) \neq 1$ ) M1  
 $\frac{dy}{dx} = \frac{(5 - 4x^2) \times f(x) - (3 - 2x^2) \times g(x)}{(5 - 4x^2)^2}$   
(either  $f(x) = -4x$  or  $g(x) = -8x$ ) A1  
 $\frac{dy}{dx} = \frac{4x}{(5 - 4x^2)^2}$  (c.a.o.) A1
6. (a) (i)  $\int \sin(x/4) dx = k \times \cos(x/4) + c$  ( $k = -1, 4, -4, -1/4$ ) M1  
 $\int \sin(x/4) dx = -4 \times \cos(x/4) + c$  A1
- (ii)  $\int e^{2x/3} dx = k \times e^{2x/3} + c$  ( $k = 1, 2/3, 3/2$ ) M1  
 $\int e^{2x/3} dx = 3/2 \times e^{2x/3} + c$  A1
- (iii)  $\int \frac{7}{8x - 2} dx = k \times 7 \times \ln|8x - 2| + c$  ( $k = 1, 8, 1/8$ ) M1  
 $\int \frac{7}{8x - 2} dx = 1/8 \times 7 \times \ln|8x - 2| + c$  A1

**Note: The omission of the constant of integration is only penalised once.**

$$(b) \int (5x + 4)^{-1/2} dx = k \times \frac{(5x + 4)^{1/2}}{1/2} \quad (k = 1, 5, 1/5) \quad \text{M1}$$

$$\int_1^9 3 \times (5x + 4)^{-1/2} dx = \left[ 3 \times \frac{1}{5} \times \frac{(5x + 4)^{1/2}}{1/2} \right]_1^9 \quad \text{A1}$$

A correct method for substitution of limits in an expression of the form  $m \times (5x + 4)^{1/2}$  M1

$$\int_1^9 3 \times (5x + 4)^{-1/2} dx = \frac{42}{5} - \frac{18}{5} = \frac{24}{5} = 4.8$$

(f.t. only for solutions of 24 and 120 from  $k = 1, 5$  respectively) A1

**Note: Answer only with no working shown earns 0 marks**

7. (a) Trying to solve either  $4x - 5 \geq 3$  or  $4x - 5 \leq -3$  M1

$$4x - 5 \geq 3 \Rightarrow x \geq 2$$

$$4x - 5 \leq -3 \Rightarrow x \leq 1/2 \quad (\text{solving both inequalities correctly}) \quad \text{A1}$$

$$\text{Required range: } x \leq 1/2 \text{ or } x \geq 2 \quad (\text{f.t. one slip}) \quad \text{A1}$$

**Alternative mark scheme**


$$(4x - 5)^2 \geq 9 \quad (\text{forming and trying to solve quadratic}) \quad \text{M1}$$

$$\text{Critical values } x = 1/2 \text{ and } x = 2 \quad \text{A1}$$

$$\text{Required range: } x \leq 1/2 \text{ or } x \geq 2 \quad (\text{f.t. one slip}) \quad \text{A1}$$

(b)  $(3|x| + 1)^{1/3} = 4 \Rightarrow 3|x| + 1 = 4^3$  M1  
 $x = \pm 21$  A1

8. (a)




Correct shape, including the fact that the  $x$ -axis is an asymptote for

$$y = f(x) \text{ at } -\infty \quad \text{B1}$$

$$y = f(x) \text{ cuts } y\text{-axis at } (0, 1) \quad \text{B1}$$

(b) (i)



Correct shape, including the fact that  $y = -4$  is an asymptote for  $y = f(3x) - 4$  at  $-\infty$  B1

(ii)  $y = f(3x) - 4$  at cuts  $y$ -axis at  $(0, -3)$  B1

(iii)  $e^{3x} = 4 \Rightarrow 3x = \ln 4$  M1  
 $x = 0.462$  A1

**Note: Answer only with no working shown earns M0 A0**

9. (a)  $y = 3 - \frac{1}{\sqrt{x-2}} \Rightarrow 3 \pm y = \pm \frac{1}{\sqrt{x-2}}$  (separating variables) M1  
 $x - 2 = \frac{1}{(3 \pm y)^2}$  or  $\frac{1}{(y \pm 3)^2}$  m1  
 $x = 2 + \frac{1}{(3 - y)^2}$  (c.a.o.) A1  
 $f^{-1}(x) = 2 + \frac{1}{(3 - x)^2}$  (f.t. one slip) A1

(b)  $D(f^{-1}) = [2.5, 3)$   
 $[2.5$  B1  
 $3)$  B1

10. (a)  $R(f) = [3 + k, \infty)$  B1
- (b)  $3 + k \geq -2$  M1  
 $k \geq -5$  ( $\Rightarrow$  least value of  $k$  is  $-5$ )  
(f.t. candidate's  $R(f)$  provided it is of form  $[a, \infty)$  A1
- (c) (i)  $gf(x) = (3x + k)^2 - 6$  B1
- (ii)  $(3 \times 2 + k)^2 - 6 = 3$   
(substituting 2 for  $x$  in candidate's expression for  $gf(x)$   
and putting equal to 3) M1  
Either  $k^2 + 12k + 27 = 0$  or  $(6 + k)^2 = 9$  (c.a.o.) A1  
 $k = -3, -9$  (f.t. candidate's quadratic in  $k$ ) A1  
 $k = -3$  (c.a.o.) A1